

Three- and Four-Body Scattering Calculations including the Coulomb Force

A. Deltuva

Centro de Física Nuclear da Universidade de Lisboa, P-1649-003 Lisboa, Portugal

Abstract. The method of screening and renormalization for including the Coulomb interaction in the framework of momentum-space integral equations is applied to the three- and four-body nuclear reactions. The Coulomb effect on the observables and the ability of the present nuclear potential models to describe the experimental data is discussed.

The Coulomb interaction, due to its long range, does not satisfy the mathematical properties required for the formulation of the standard scattering theory. However, since in nature the Coulomb potential is always screened, one could expect that the physical observables become insensitive to the screening provided it takes place at sufficiently large distances R and, therefore, the $R \rightarrow \infty$ limit should correspond to the proper Coulomb. This was proved by Taylor [1] in the context of the two-particle system: though the on-shell screened Coulomb transition matrix diverges in the $R \rightarrow \infty$ limit, after renormalization by (an equally) diverging phase factor it converges as a distribution to the well known proper Coulomb amplitude and therefore yields identical results for the physical observables. A similar renormalization relates screened and proper Coulomb wave functions [2].

The method of screening and renormalization can be used for the systems with more particles [3], albeit with some limitations. Here we briefly recall the procedure which is described in detail in ref. [4]. In the transition operators derived from nuclear plus screened Coulomb potentials one has to isolate the diverging screened Coulomb contributions in the form of a two-body on-shell transition matrix and two-body wave function with known renormalization properties. This can be achieved using the two-potential formalism as long as in the initial/final states there are no more than two charged bodies (clusters). At the same time this procedure separates long-range and Coulomb-distorted short-range parts of the transition amplitude, the former being the two-body on-shell transition matrix derived from the screened Coulomb potential between the centers of mass (c.m.) of the two charged bodies that is present in the elastic scattering only. After renormalization this contribution converges towards its $R \rightarrow \infty$ limit very slowly but the result, the pure Coulomb amplitude of two-body nature, is known

analytically. The remaining part of the elastic scattering amplitude as well as the amplitudes for transfer and breakup are complicated short-range operators that are externally distorted by Coulomb. However, due to their short-range nature, convergence with R after the renormalization by the corresponding phase factors is fast and, therefore, they are calculated numerically at finite R using the standard scattering theory and making sure that R is large enough for the convergence of the results. We solve Faddeev-like Alt, Grassberger, and Sandhas (AGS) equations for three- and four-particle scattering [5, 6] using the momentum-space partial-wave basis as described in detail in refs. [7, 8, 9] for three- and four-nucleon scattering without the Coulomb force. However, the screened Coulomb interaction, due to its longer range, compared to the nuclear interaction, brings additional difficulties: quasisingular nature of the potential and slow convergence of the partial-wave expansion. The right choice of the screening is essential in resolving those difficulties. The convergence of the partial-wave expansion with our new screening function [4] is fast enough and thereby allows us to avoid the approximations used in the previous implementations [10, 11] of the screening and renormalization approach and obtain reliable results.

The most important criterion for the reliability of the screening and renormalization method is the convergence of the observables with the screening radius R used to calculate the Coulomb-distorted short-range part of the amplitudes. Numerous examples can be found in refs. [4, 12, 13]. In most cases the convergence is impressively fast and only becomes slower for the observables at very low energies. Furthermore, as demonstrated in ref. [14], our results for p - d elastic scattering agree well over a wide energy range with those of ref. [15] obtained from the variational solution of the three-nucleon Schrödinger equation in configuration space with the inclusion of an *unscreened* Coulomb potential and imposing the proper Coulomb boundary conditions explicitly.

The present method was used to study three-nucleon hadronic and electromagnetic (e.m.) reactions in refs. [4, 16, 17, 18]. Furthermore, it was applied to the nuclear reactions dominated by three-body degrees of freedom like $\alpha + d$ [12], $d + {}^{12}\text{C}$, and $p + {}^{11}\text{Be}$ [19, 20]. Finally, in refs. [9, 13, 21] all elastic and transfer four-nucleon reactions below three-body breakup threshold have been studied. The importance of the Coulomb at low energies is demonstrated in Fig. 1 for elastic d - α scattering. It may be very strong at all energies in p - d breakup and three-body e.m. disintegration of ${}^3\text{He}$ in kinematical regimes with low relative pp energy where the Coulomb repulsion converts the cross section peak obtained in the absence of Coulomb into a minimum as can be seen in the experimental data as well [16, 18]. However, even after the inclusion of the Coulomb interaction and the three-nucleon force some discrepancies between experiment and theory like the space star anomaly in p - d breakup [4, 18] and the A_y -puzzle in p - d [4, 15] and p - ${}^3\text{He}$ [13, 24] elastic scattering still persist. Furthermore, A_y is described quite well in the n - ${}^3\text{He}$ and p - ${}^3\text{H}$ elastic scattering but not in the $p + {}^3\text{H} \rightarrow n + {}^3\text{He}$ transfer reaction. A very strong Coulomb effect manifests itself in the α - d breakup where the shift of αp P -wave resonance position leads to the corresponding shifts of the differential cross section peaks as shown in Fig. 2. In addition, Figs. 1 and 2 as well as the results of ref. [12] demonstrate the su-




Figure 1. Differential cross section and deuteron vector analyzing power iT_{11} of d - α elastic scattering at $E_d = 4.81$ MeV. Results derived from the N - α potential that is attractive in S-wave and supports Pauli-forbidden bound state which is projected out are shown as solid (dashed) curves with (without) Coulomb. The results including Coulomb but with local repulsive N - α S-wave potential are given by dotted curves. The experimental data are from refs. [22, 23]




Figure 2. Differential cross section of α - d breakup at $E_\alpha = 15$ MeV in selected kinematical configurations. Curves as in Fig. 1. The experimental data are from ref. [22, 23]

priority of the attractive N - α S-wave potentials supporting a Pauli-forbidden bound state that is projected out over the local repulsive S-wave potentials which, because of their simplicity, are very often used in the configuration space calculations of resonances and e.m. reactions.

In conclusion, the Coulomb interaction between the charged particles was included in few-body scattering calculations using the old idea of screening and renormalization [1] but with novel practical realization that avoids all the approximations of the previous works [10, 11] and yields fully converged results.

Acknowledgement. This work has been performed in collaboration with A. C. Fonseca and P. U. Sauer. The author is supported by the Fundação para a Ciência e a Tecnologia (FCT) grant SFRH/BPD/34628/2007.

References

1. Taylor, J. R.: Nuovo Cim. **B23**, 313 (1974)
2. Gorshkov, V. G.: Sov. Phys.-JETP **13**, 1037 (1961)
3. Alt, E. O., Sandhas, W.: Phys. Rev. **C21**, 1733 (1980)
4. Deltuva, A., Fonseca, A. C., Sauer, P. U.: Phys. Rev. **C71**, 054005 (2005); Phys. Rev. Lett. **95**, 092301; Phys. Rev. **C72**, 054004 (2005); Phys. Rev. **C73**, 057001 (2006); Annu. Rev. Nucl. Sci. **58**, 27 (2008)
5. Alt, E. O., Grassberger, P., Sandhas, W.: Nucl. Phys. **B2**, 167 (1967)
6. Grassberger, P., Sandhas, W.: Nucl. Phys. **B2**, 181 (1967); Alt, E. O., Grassberger, P., Sandhas, W.: JINR report No. E4-6688 (1972)
7. Chmielewski, K., et al.: Phys. Rev. **C67**, 014002 (2003)
8. Deltuva, A., Chmielewski, K., Sauer, P. U.: Phys. Rev. **C67**, 034001 (2003)
9. Deltuva, A., Fonseca, A. C.: Phys. Rev. **C75**, 014005 (2007)
10. Alt, E. O., Rauh, M.: Few-Body Systems **17**, 121 (1994)
11. Alt, E. O., et al.: Phys. Rev. **C65**, 064613 (2002)
12. Deltuva, A.: Phys. Rev. **C74**, 064001 (2006)
13. Deltuva, A., Fonseca, A. C.: Phys. Rev. Lett. **98**, 162502 (2007); Phys. Rev. **C76**, 021001 (2007)
14. Deltuva, A., et al.: Phys. Rev. **C71**, 064003 (2005)
15. Kievsky, A., Viviani, M., Rosati, S.: Phys. Rev. **C64**, 024002 (2001)
16. Kistryn, S., et al.: Phys. Lett. **B641**, 23 (2006)
17. Deltuva, A., Fonseca, A. C., Sauer, P. U.: Nucl. Phys. **A790**, 344c (2007)
18. Sagara, K., et al.: Nucl. Phys. **A790**, 348c (2007)
19. Deltuva, A., et al.: Phys. Rev. **C76**, 064602 (2007)
20. Crespo, R., et al.: Phys. Rev. **C76**, 014620 (2007)
21. Deltuva, A., Fonseca, A. C., Sauer, P. U.: Phys. Lett. **B660**, 471 (2008)
22. Bruno, M., et al.: Lett. Nuovo Cim. **27**, 265 (1980)
23. Grüebler, W., et al.: Nucl. Phys. **A134**, 686 (1969)
24. Viviani, M., et al.: Phys. Rev. Lett. **86**, 3739 (2001)
25. Koersner, I., et al.: Nucl. Phys. **A286**, 431 (1977)